

4. If $a \in \mathbb{R}$ satisfies $a \cdot a = a$, prove that either $a = 0$ or $a = 1$.
5. If $a \neq 0$ and $b \neq 0$, show that $1/(ab) = (1/a)(1/b)$.
6. Use the argument in the proof of Theorem 2.1.4 to show that there does not exist a rational number s such that $s^2 = 6$.
7. Modify the proof of Theorem 2.1.4 to show that there does not exist a rational number t such that $t^2 = 3$.
8. ✓(a) Show that if x, y are rational numbers, then $x + y$ and xy are rational numbers.
(b) Prove that if x is a rational number and y is an irrational number, then $x + y$ is an irrational number. If, in addition, $x \neq 0$, then show that xy is an irrational number.
9. Let $K := \{s + t\sqrt{2} : s, t \in \mathbb{Q}\}$. Show that K satisfies the following:
 - (a) If $x_1, x_2 \in K$, then $x_1 + x_2 \in K$ and $x_1 x_2 \in K$.
 - (b) If $x \neq 0$ and $x \in K$, then $1/x \in K$.
 (Thus the set K is a *subfield* of \mathbb{R} . With the order inherited from \mathbb{R} , the set K is an ordered field that lies between \mathbb{Q} and \mathbb{R} .)
- ✓10. (a) If $a < b$ and $c \leq d$, prove that $a + c < b + d$.
(b) If $0 < a < b$ and $0 \leq c \leq d$, prove that $0 \leq ac \leq bd$.
11. (a) Show that if $a > 0$, then $1/a > 0$ and $1/(1/a) = a$.
(b) Show that if $a < b$, then $a < \frac{1}{2}(a + b) < b$.
12. Let a, b, c, d be numbers satisfying $0 < a < b$ and $c < d < 0$. Give an example where $ac < bd$, and one where $bd < ac$.
13. If $a, b \in \mathbb{R}$, show that $a^2 + b^2 = 0$ if and only if $a = 0$ and $b = 0$.
14. If $0 \leq a < b$, show that $a^2 \leq ab < b^2$. Show by example that it does *not* follow that $a^2 < ab < b^2$.
15. If $0 < a < b$, show that (a) $a < \sqrt{ab} < b$, and (b) $1/b < 1/a$.
- ✓16. Find all real numbers x that satisfy the following inequalities.

(a) $x^2 > 3x + 4$,	(b) $1 < x^2 < 4$,
(c) $1/x < x$,	(d) $1/x < x^2$.
- ✓17. Prove the following form of Theorem 2.1.9: If $a \in \mathbb{R}$ is such that $0 \leq a \leq \varepsilon$ for every $\varepsilon > 0$, then $a = 0$.
18. Let $a, b \in \mathbb{R}$, and suppose that for every $\varepsilon > 0$ we have $a \leq b + \varepsilon$. Show that $a \leq b$.
19. Prove that $[\frac{1}{2}(a + b)]^2 \leq \frac{1}{2}(a^2 + b^2)$ for all $a, b \in \mathbb{R}$. Show that equality holds if and only if $a = b$.
20. (a) If $0 < c < 1$, show that $0 < c^2 < c < 1$.
(b) If $1 < c$, show that $1 < c < c^2$.
21. (a) Prove there is no $n \in \mathbb{N}$ such that $0 < n < 1$. (Use the Well-Ordering Property of \mathbb{N} .)
(b) Prove that no natural number can be both even and odd.
22. (a) If $c > 1$, show that $c^n \geq c$ for all $n \in \mathbb{N}$, and that $c^n > c$ for $n > 1$.
(b) If $0 < c < 1$, show that $c^n \leq c$ for all $n \in \mathbb{N}$, and that $c^n < c$ for $n > 1$.
23. If $a > 0, b > 0$, and $n \in \mathbb{N}$, show that $a < b$ if and only if $a^n < b^n$. [Hint: Use Mathematical Induction.]
24. (a) If $c > 1$ and $m, n \in \mathbb{N}$, show that $c^m > c^n$ if and only if $m > n$.
(b) If $0 < c < 1$ and $m, n \in \mathbb{N}$, show that $c^m < c^n$ if and only if $m > n$.
25. Assuming the existence of roots, show that if $c > 1$, then $c^{1/m} < c^{1/n}$ if and only if $m > n$.
26. Use Mathematical Induction to show that if $a \in \mathbb{R}$ and $m, n \in \mathbb{N}$, then $a^{m+n} = a^m a^n$ and $(a^m)^n = a^{mn}$.